



SPARC Workshop SHARP2016

Diagnosis of the Upper Troposphere-Stratosphere exchanges using CO₂ tracer gas

Dr. Mohamadou Diallo¹, Prof. Bernard Legras², Dr. Eric Ray³, Prof. Alain Chédin⁴, Prof. Andreas Engel⁵, Junior Prof. Juan Antonio Añel Cabanelas⁶, Prof. Laura de la Torre Ramos¹

¹ EPhysLab, Faculty of Sciences, University of Vigo, Ourense, Spain.

² Laboratoire de Météorologie Dynamique, UMR8539, IPSL, UPMC/ENS/CNRS/Ecole Polytechnique, Paris, France.

³ Chemical Sciences Division, ESRL,NOAA and CIRES,University of Colorado, Bould

mdiallo@lmd.ens.fr

Currently, most of our knowledge of the upper-tropospheric-stratospheric global distribution of the carbon dioxide (CO₂), is limited by the uncertainty in the sparseness of in situ measurements and in transport and mixing processes. With a strengthening shallow branch of the Brewer-Dobson circulation (BDC), the CO₂ and other halocarbons increases is expected in the UT/LS and can change the stratospheric radiative forcing.

This study uses the estimated mean AoA and the inert feature of the CO₂ over the 2000-2010 period to investigate the stratospheric transport and mixing processes and the possible increase of the stratospheric CO₂ due to the BDC changes. The monthly mean CO₂ and AoA reconstructed from the backward trajectories guided by ERA-Interim re-analyses is compared with those from stratospheric balloons, SOLVE and CONTRAIL campaigns and ACE-FTS satellite retrieval CO₂. The inter-comparison between the in situ observations and reconstructions revealed a fairly good agreement. That demonstrates the potentiality of using lagrangian reconstruction in order to better understand the stratospheric transport and mixing processes. The zonal distribution exhibits a high CO₂ in the tropical stratosphere due to the tropical upwelling of BDC and to the efficiency of Rossby wave strengthening the fast quasi-horizontal mixing. In winter and spring, the tropical pipe is confined in the tropical band while it is spread during summer and autumn permitting high CO₂ to reach the high stratosphere. The seasonal variations of the mean AoA and CO₂ anomalies are dominated by the seasonal variation of the mixing barriers such as the subtropical barrier, which isolates the tropical pipe from surf zone in the overworld, and the tropopause barrier, which limits cross-tropopause mixing between the tropical tropopause layer and the mid-latitude lowermost stratosphere in winter. The variation of the barriers permeability seasonally modulates the efficiency of the UT/LS exchanges.